

Water purification using Iodine crystals.

1 oz. glass jar (must have glass or bakelite stopper otherwise the iodine will react with the plastic or metal stopper and destroy it.) Actually 0.1 gram is adequate for the job, but using a larger amount of iodine creates a saturated solution much quicker.

Put in 1 oz. (1 tablespoon or 3 teaspoons) of water (at least room temperature, body temperature preferred).

Close stopper and shake for several minutes. You now have a saturated solution. A saturated solution is when as much solid has dissolved in a liquid as it can.

Carefully pour off 10ml (10cc, 2 teaspoons) of the saturated solution.

REMEMBER, the iodine crystals are VERY TOXIC! The reason that adding more water than needed is suggested is so that you need not tip the bottle over too far thus spilling some crystals.

Add the 10ml (2 teaspoons) of saturated solution to 1 liter (1.06 quart) of water.

Let stand at least 15 minutes at 77 degrees F. or higher. Make sure all of the interior surface including lid get treated.

How to use ionic iodine to counter radioactive iodine contamination.

KI (potassium Iodide) (reagent grade is best)

one pound of KI = 3,500 doses = treatment for 35 adults or children for 100 days.

Purchase crystalline or granular KI (reagent grade).

Fill a glass jar 60% full of KI (2 oz. of KI in a 2 Oz. bottle *dark glass*)

Fill with water (90%)

Shake for two minutes.

Some of the KI should remain undissolved (proof of saturated solution!)

1 drop of saturated solution = 28 to 36 mg. of KI

130 mg KI per day (4 drops) for Adults (FDA recommended daily dose), 65 mg per day (2 drops) for infants under 1 year.

Radioactive iodine has a 8 day half-life, so 80 days = .1% (1/1,000)

Period of dosage, start dosage beginning of first warning of a radioactive iodine release from nuclear reactor accident or nuclear bomb detonation, continue until at least 10 days beyond most recent release of radioactive Iodine 131 is recommended minimum, 100 days would be the most conservative method of treatment. Having only doses for a day or two is VASTLY better than none at all.